

The Generalization of the Keldysh Theorem

MILUTIN R. DOSTANIĆ

*Institute for Applied Mathematics and Electronics,
Kneza Miloša 37, Belgrade, Yugoslavia*

Submitted by James S. Howland

Received March 21, 1989

In this paper we generalize the theorem by M. V. Keldysh on completeness of a system root vectors of a weakly perturbed m -sectorial compact operator with nontrivial kernel. © 1991 Academic Press, Inc.

1. INTRODUCTION AND NOTATION

In this paper we generalize the theorem by M. V. Keldysh [4] on completeness of a system root vectors of weakly perturbed injective, compact selfadjoint operators H to the case of weakly perturbed m -sectorial compact operator with nontrivial kernel.

Let \mathcal{H} be a separable Hilbert space over \mathbb{C} , and $A \in \mathcal{L}(\mathcal{H})$ be a compact operator. The eigenvalues of the compact nonnegative selfadjoint operator $(A^*A)^{1/2}$ arranged in decreasing order and repeated according to multiplicity, form a sequence s_1, s_2, s_3, \dots approaching zero. Following V. Neumann Schatten [1, Sect. XI, 9] we say that A belongs to the class C_p ($0 < p < \infty$) provided $\sum_j s_j^p < \infty$. The set C_p is a two sided ideal in $\mathcal{L}(\mathcal{H})$. The number $\lambda \in \mathbb{C}$ is called a characteristic value of A if there exists $x \neq 0$ ($x \in \mathcal{H}$) such that $(I - \lambda A)x = 0$, of course $\lambda \neq 0$ and x is an eigenvector of A for the eigenvalue λ^{-1} . Further $x \neq 0$ is a generalized eigenvector (for the characteristic value λ) if for some $k \geq 1$

$$(I - \lambda A)^k x = 0.$$

Let $\overline{\mathcal{R}(A)}$ denote the closure of the range of A . In [3], the generalization of the Keldysh theorem to the case of weakly perturbed selfadjoint operators with nontrivial kernel under some additional assumption is given.

2. MAIN RESULT

THEOREM. Let $B = A(I + S)$ where S is a compact operator $-1 \in \rho(S)$, and $\text{Ker } A \subset \text{Ker } S$. If the operator $A \in C_p$ ($0 < p < \infty$) is such that the values of quadratic form (Af, f) lie in the sector

$$\Omega = \left\{ \lambda : |\arg \lambda| \leq \theta < \frac{\pi}{2\rho'} \right\}, \quad \text{where } \rho' > \max\{1, p\},$$

then:

(a) For all ε , $0 < \varepsilon < \pi/2 - \theta$, all characteristic values of operator B , except finitely many, lie in the sector $F_\varepsilon = \{\lambda : |\arg \lambda| < \theta + \varepsilon\}$.

(b) The system of generalized eigenvectors of the operator B is complete in $\overline{\mathcal{R}(A)}$.

Before we prove this theorem first we show the following

LEMMA. Under the assumptions given in the previous theorem

$$\lim_{\substack{\lambda \rightarrow \infty \\ \lambda \in F_\varepsilon^c}} \|S(I - \lambda A)^{-1}\| = 0, \quad (1)$$

where $F_\varepsilon^c = \mathbb{C} \setminus F_\varepsilon$ and convergence is uniform on F_ε^c .

Proof. Because of given assumptions $\{(Af, f) : f \in \mathcal{H}\} \subset \Omega$ so [2] holds $\|(A - \mu I)^{-1}\| \leq 1/d(\mu, W_A)$ ($\mu \notin W_A$), where W_A is the closure of set $\{(Af, f) : \|f\| = 1\}$ and $d(\mu, W_A)$ is the distance between point μ and set W_A .

If $\lambda \in F_\varepsilon^c$ for $\mu = \lambda^{-1}$ we have $\mu \in F_\varepsilon^c$ and one has

$$\begin{aligned} \|(I - \lambda A)^{-1}\| &\leq \frac{1}{\sin \varepsilon} \\ \|(I - \bar{\lambda} A^*)^{-1}\| &\leq \frac{1}{\sin \varepsilon}. \end{aligned} \quad (2)$$

Since the operator S is compact it can be represented in the form

$$S = \sum_k \mu_k(\cdot, f_k) g_k \quad (\mu_1 = \|S\| \geq \mu_2 \geq \mu_3 \geq \dots).$$

The system of vectors $\{g_k\}$ and $\{f_k\}$ are orthonormal systems, so that

$$S^* g_k = \mu_k f_k. \quad (3)$$

The assumptions of the Theorem imply that $\overline{\mathcal{R}(A^*)} \supset \overline{\mathcal{R}(S^*)}$. This inclusion and (3) imply that $f_k \in \overline{\mathcal{R}(A^*)}$ for every $k \in N$. Since S can be uniformly approximated by finite rank operators, to prove the Lemma it is enough to show (1) for an operator S of the form

$$S = \sum_{k=1}^n \mu_k(\cdot, f_k) g_k.$$

But, in order to show that it is enough to consider S of the form

$$S = \mu_k(\cdot, f_k) g_k \quad (\forall k \in N).$$

For $S = \mu_k(\cdot, f_k) g_k$ it is easy to see that

$$\|S(I - \lambda A)^{-1}\| \leq \|(I - \lambda A^*)^{-1} f_k\| \cdot \|S\| \quad (\text{for } \lambda \in F_\varepsilon^c).$$

So, it is enough to prove that

$$\lim_{\substack{\lambda \rightarrow \infty \\ \lambda \in F_\varepsilon^c}} \|(I - \lambda A^*)^{-1} f_k\| = 0 \quad (\forall k \in N) \text{ uniformly on } F_\varepsilon^c. \quad (4)$$

Now, one can apply the theorem of Lidsky [5].

Let $h \in \overline{\mathcal{R}(A^*)}$. Thus for every fixed α (such that $\max\{1, p\} \leq \alpha < \pi/2\theta$) the Fourier series of the vector h with respect to the system of main vectors $\{G_s\}$ (in the sense of [5]) of A^* is summable by Abel's method of order α . So

$$h = \lim_{t \rightarrow 0+} \sum_{v=1}^{\infty} \left(\sum_{s=N_v+1}^{N_{v+1}} C_s(t) G_s \right). \quad (5)$$

(The above is the fundamental theorem in [5]. The coefficients $C_s(t)$ are defined in [5].) The series in (5) is convergent in \mathcal{H} for every fixed $t > 0$.

From (5) it follows that the system of vectors is complete in $\overline{\mathcal{R}(A^*)}$. Since $f_k \in \overline{\mathcal{R}(A^*)}$, the lemma follows from (2), (4), (5) and

$$\lim_{\substack{\lambda \rightarrow \infty \\ \lambda \in F_\varepsilon^c}} (I - \lambda A^*)^{-1} G_s = 0 \quad (\forall s \in N) \text{ uniformly on } F_\varepsilon^c. \quad (6)$$

The system of vectors $\{G_s\}$ is the union of Jordan's chains of main vectors of operator A^* [5] which correspond to different eigenvalues of A^* . So, it is enough to prove (6) for one such chain.

Suppose then, that λ_0 is an eigenvalue of operator A^* , G_0 is the corresponding eigenvector, and G_1, G_2, \dots, G_k is a Jordan chain of vectors. (There can be only finitely many vectors G_i because A^* is compact).

So, $A^*G_0 = \lambda_0 G_0$, $A^*G_1 = \lambda_0 G_1 + G_0$, ..., $A^*G_k = \lambda_0 G_k + G_{k-1}$. It can be established directly that

$$(I - \bar{\lambda}A^*)^{-1} G_0 = \frac{G_0}{1 - \bar{\lambda}\lambda_0} \quad (7)$$

$$\begin{aligned} (I - \bar{\lambda}A^*)^{-1} G_1 &= \frac{\bar{\lambda}}{1 - \bar{\lambda}\lambda_0} (I - \bar{\lambda}A^*)^{-1} G_0 + \frac{G_1}{1 - \bar{\lambda}\lambda_0} \\ &\vdots \\ (I - \bar{\lambda}A^*)^{-1} G_k &= \frac{\bar{\lambda}}{1 - \bar{\lambda}\lambda_0} (I - \bar{\lambda}A^*)^{-1} G_{k-1} + \frac{G_k}{1 - \bar{\lambda}\lambda_0}. \end{aligned} \quad (8)$$

Now (6) can be obtained directly from (7) and (8). So the proof of the Lemma is completed.

Notice that

$$\lim_{\substack{\lambda \rightarrow \infty \\ \lambda \in F_\varepsilon^c}} \|T(I - \lambda A)^{-1}\| = 0 \quad (9)$$

because of the Lemma, where $T = S(I + S)^{-1}$ and the convergence is uniform on F_ε^c .

Proof of the Theorem. The proof of the Theorem is now based on an idea of Keldysh [4].

(a) From $B = A(I + S)$ it follows

$$\begin{aligned} (I - \lambda B)^{-1} &= (I + S)^{-1} (I - T(I - \lambda A)^{-1})^{-1} (I - \lambda A)^{-1}, \\ \text{where } T &= S(I + S)^{-1}. \end{aligned} \quad (10)$$

Because of the Lemma, there exists $r > 0$ so that for $|\lambda| > r$ and $\lambda \in F_\varepsilon^c$ one has

$$\|T(I - \lambda A)^{-1}\| \leq a < 1 \quad (a = \text{const}).$$

So the formula (10) is correct and implies part (a) of the result and estimate

$$\|(I - \lambda B)^{-1}\| \leq \frac{\|(I + S)^{-1}\|}{(1 - a) \sin \varepsilon} \quad (11)$$

for $\lambda \in F_\varepsilon^c$ and $|\lambda| > r$.

(b) Let \mathcal{S} be the closed span of all generalized eigenvectors of the operator B .

Clearly $\mathcal{S} \subset \overline{\mathcal{R}(B)}$. Let P be an orthogonal projector of space \mathcal{H} on \mathcal{S} and $Q = I - P$. Thus $B_1 = QBQ$ is a Volterra operator [2], so the operator function $(I - \lambda B_1)^{-1}$ is an entire function and [2] one has

$$(I - \lambda B_1)^{-1} = Q(I - \lambda B)^{-1} Q + P. \quad (12)$$

Chose $\varepsilon > 0$ so that $\theta + \varepsilon < \pi/2p$. Since $A \in C_p$ one has $B \in C_p$ and $B_1 \in C_p$ and [2]

$$\ln \|(I - \lambda B_1)^{-1}\| = O(|\lambda|^p). \quad (13)$$

From (11), (12), and (13) it follows that the function $(I - \lambda B_1)^{-1}$ is bounded on boundary of sector F_ε and since $2(\theta + \varepsilon) < \pi/p$, by the theorem of Phragmen, Lindelof, and Liouville, follows

$$B_1 = 0 \quad \text{or} \quad QBQ = 0.$$

Then, we have $QB = 0$ or $B^*Q = 0$, i.e., $(I + S^*)A^*Q = 0$. Thus

$$A^*Q = 0. \quad (14)$$

From (14) follows $Q\mathcal{H} \subset \text{Ker } A^*$, or $(Q\mathcal{H})^\perp \supset (\text{Ker } A^*)^\perp$. Since $(Q\mathcal{H})^\perp = P\mathcal{H} = \mathcal{S}$ and $(\text{Ker } A^*)^\perp = \mathcal{R}(A)$ we have

$$\overline{\mathcal{R}(A)} \subset \mathcal{S}.$$

But, because of the assumption $-1 \in \rho(S)$, one has $\overline{\mathcal{R}(B)} = \mathcal{R}(B)$ so that

$$\overline{\mathcal{R}(B)} \subset \mathcal{S}. \quad (15)$$

Since $\mathcal{S} \subset \overline{\mathcal{R}(B)}$, (15) implies $\mathcal{S} = \overline{\mathcal{R}(B)}$ i.e.,

$$\mathcal{S} = \overline{\mathcal{R}(A)}.$$

This completes the proof of the Theorem.

Remark 1. If $p \geq 1$ the assumption $\text{Ker } A \subset \text{Ker } S$ is not necessary. In that case, by Lidsky's theorem [2] $\text{Ker } A = \{0\}$ and condition $\text{Ker } A = \text{Ker } S$ is satisfied.

Remark 2. If $p \geq 1$ the theorem can be formulated in the following way: If T is a compact operator and $A \in C_p$ is such that

$$\{(Af, f) : f \in \mathcal{H}\} \subset \Omega = \left\{ \lambda : |\arg \lambda| \leq \theta < \frac{\pi}{2p} \right\}$$

then the system of eigenvectors and associated vectors [2] of the linear pencil $L(\lambda) = I - T - \lambda A$ is complete in \mathcal{H} .

EXAMPLE. The operator L generated by the differential expression

$$l(y) = -y'' + p_1(x) y' + p_2(x) y$$

and boundary condition is m -sectorial if for example

$$y(0) = 0, \quad y(\pi) = 0$$

$$\max_{0 \leq x \leq \pi} |p_1(x)|^2 \leq \min_{0 \leq x \leq \pi} |p_2(x)|,$$

where p_1, p_2 are real continuous functions on $[0, \pi]$. Suppose that the equation $Ly = 0$ has only the trivial solution $y = 0$.

If $K(\cdot, \cdot)$ is a continuous function on $[0, \pi] \times [0, \pi]$ so that

$$\int_0^\pi \int_0^\pi |K(x, s)|^2 dx ds < 1$$

and the boundary value problem

$$-y'' + p_1(x) y' + p_2(x) y = \lambda y + \lambda \int_0^\pi K(x, s) y(s) ds$$

$$y(0) = y(\pi) = 0$$

has simple eigenvalues, then the corresponding system of eigenvectors is complete in $L^2(0, \pi)$.

The proof follows directly by applying the theorem to the operator $A = L^{-1}(I + S)$ where $S : L^2(0, \pi) \rightarrow L^2(0, \pi)$ is the linear operator defined by

$$Sf(x) = \int_0^\pi K(x, s) f(s) ds.$$

REFERENCES

1. N. DUNFORD AND J. T. SCHWARTZ, "Linear Operators," Part II, Interscience, New York/London, 1963.
2. I. C. GOHBERG AND M. G. KREIN, Introduction to the theory of linear nonselfadjoint operators, in "Translations of Mathematical Monographs," Vol. 18, Amer. Math. Soc., Providence, RI, 1969.
3. P. HESS, On the relative completeness of the generalized eigenvectors of elliptic eigenvalue problems with indefinite weight functions, *Math. Ann.* **270** (1985), 467-475.
4. M. V. KELDYSH, On the characteristic values and characteristic functions of certain classes of nonselfadjoint equations, *Dokl. Akad. Nauk USSR* **77** (1951), 11-14.
5. V. B. LIDSKY, On summability of series on main vectors for nonselfadjoint operators, *Trudy Moskov. Mat. Obshch.* **11** (1962), 3-35.